


Certificate of Analysis			
Vector Name	PCC1BAC	Catalog No.	V008675
Project/Lot No.	C065NJTPG0-1/PD89096	Strain	EPI300
Quantity	200ug	Resistance	Chloramphenicol
QC Results			
Test Items	Specifications	Results	
Appearance	Clear, no visible particles	Pass	
Concentration	On request, default	Pass	
A260/280	1.8 - 2.0	Pass	
A260/230	>2.0	Pass	
Supercoiled content	No request	N/A	
Residual RNA	Not visible upon electrophoresis	Pass	
Genomic DNA	Not visible upon electrophoresis	Pass	
Restriction Analysis	Conforms to reference	Pass	
Bioburden assay	No growth on agar plate after 24 hours	N/A	
Endotoxin Test	Verified, (Endo-Free Preps Only)	N/A	
Label	Correct and white	Pass	
Sequencing	Correct	Pass	
Comments	-	-	
Restriction Digestion Map			
	1	2	M
	<p>Lane 1: Plasmid DNA</p> <p>Lane 2: Plasmid digested by EcoRI_AgeI (3495/2521/1230/893 bp in theory)</p> <p>Lane M: KB Ladder</p>		

Certified by: Chou Fang
Date: Oct/18/2024
Valid until: Oct/18/2026



Note

BAC and fosmid clones are highly suitable for modification by recombineering but, because they are present at low (1-2) copies per cell, the DNA is difficult to isolate in high yield and purity. To overcome this limitation vectors, e.g. pCC1BAC/pCC1FOS, have been constructed that contain the additional replication origin, *oriV*, which permits copy-number to be induced transiently when propagated in a suitable host strain, e.g. EPI300, that supplies the cognate trans-replication protein TrfA.

Protocol for EPI400/EPI300

- Add 4 ml LB media into each test tube. Inoculate each tube with bacterial culture with antibiotic at the proper concentration.
- Incubate the tubes at 37°C, shaking overnight.
- Dilute the starting culture (from step b) 1:10 into antibiotic-supplemented fresh media.
- Supplement induction solution with a ratio of 1:1000, and grow the culture at 37°C for 4 h with vigorous shaking (approx. 250 rpm).
- Isolate DNA from the induced culture cells as per the protocol provided

Vector Map

