myc-BioID2-MCS

载体编号 Vector ID	V000701
抗性 Antibiotic Resistance	Ampicillin
长度 Length	6146 bp
载体类型 Vector Type	Mammalian Expression

产品信息

复制子 Replication Origin	ori
筛选标记 Selection Marker	Neomycin (select with G418)
启动子 Promoter	CMV
克隆方法 Cloning Method	Restriction Enzyme
5' 引物 5' Primer	CMV-F
3' 引物 3' Primer	BGH-rev
培养菌株 Growth Strain(s)	DH5alpha
培养温度 Growth Temperature	37℃
备注 Comment	The myc-BioID2-MCS construct is built on the pcDNA3.1 vector backbone. This tool allows for the fusion of your protein of interest to the C-terminus of BioID2, facilitating proximity-dependent biotin identification (BioID). The multiple cloning site (MCS) is positioned downstream of the myc tag and the BioID2 tag.

质粒图谱



复溶方法

1. Centrifuge at 5,000×g for 5 min.
2. Carefully open the tube and add sterile water to dissolve the DNA: add 20 µl for 5 µg plasmid, and 100 µl for 100 µg plasmid.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than 5000×g.
5. Store the plasmid at -20 ℃.