pDisplay

载体编号 Vector ID	V000792
抗性 Antibiotic Resistance	Ampicillin
长度 Length	5324 bp
载体类型 Vector Type	Mammalian Surface Display System

产品信息

复制子 Replication Origin	ori
筛选标记 Selection Marker	Neomycin
启动子 Promoter	SV40
5' 引物 5' Primer	T7-F:TAATACGACTCACTATAGGG
融合标签 Fusion Tag	HA Tag, c-Myc Epitope Tag, PDGFR Transmembrane Domain, IgK Leader Sequence
备注 Comment	pDisplay is a mammalian expression vector designed to target recombinant proteins to the surface of mammalian cells. Displayed proteins can be analyzed for their ability to interact with known or putative ligands. Proteins of interest are targeted and anchored to the cell surface by cloning the gene of interest in frame with the vectors unique N-terminal secretion signal and the C-terminal transmembrane anchoring domain of platelet-derived growth factor receptor (PDGFR). In contrast to phage display vectors which operate exclusively in prokaryotic cells, the pDisplay vector offers the advantage of having the displayed protein of interest processed in mammalian cells. Therefore, recombinant proteins of eukaryotic origin that are expressed from pDisplay more closely resemble their native form. This may favor more accurate ligand binding interactions. In addition to the N-terminal cell surface targeting signal and C-terminal transmembrane anchoring domain, the pDisplay vector contains the following key features:• T7 promoter/priming site for in vitro transcription of sense RNA and for sequencing of inserts • Neomycin resistance marker for stable selection in mammalian cells using Geneticin • SV40 origin for replication and simple vector rescue in cell lines expressing the large T antigen (e.g., COS-1 and COS-7) • Ampicillin resistance gene for selection in E. coli

质粒图谱



复溶方法

1. Centrifuge at 5,000×g for 5 min.
2. Carefully open the tube and add sterile water to dissolve the DNA: add 20 µl for 5 µg plasmid, and 100 µl for 100 µg plasmid.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than 5000×g.
5. Store the plasmid at -20 ℃.