pLVX-shRNA1

载体编号 Vector ID	V012697
抗性 Antibiotic Resistance	Ampicillin
长度 Length	7757 bp
载体类型 Vector Type	Viral Expression & Packaging Vectors

产品信息

复制子 Replication Origin	ori
来源 Source/Author	Clontech
筛选标记 Selection Marker	Puromycin
拷贝数 Copy Number	High copy number
启动子 Promoter	mPGK
克隆方法 Cloning Method	Enzyme Cut
表达方式 Expression Method	Constiutive, Stable
备注 Comment	pLVX-shRNA1 is an HIV-1-based, lentiviral expression vector designed to express a small hair-pin RNA (shRNA) for RNA interference (RNAi) studies. Expression of your shRNA is drivenby the RNA Pol III-dependent, human U6 promoter (PU6), located just upstream of the MCS. pLVX-shRNA1 can either be used as a plasmid expression vector and transfected into cells, or it can be packaged into viral particles and transduced into cells. Lentiviral particles derived from the vector allow the expression of shRNAs in virtually any cell type, including primary cells. pLVX-shRNA1 contains all of the viral processing elements necessary for the production of replication-incompetent lentivirus, as well as elements to improve viral titer and overall vector function. The woodchuck hepatitis virus posttranscriptional regulatory element (WPRE) promotes RNA processing events and enhances nuclear export of viral RNA (1), leading to increased viral titers from packaging cells. In addition, the vector includes a Rev-response element (RRE), which further increases viral titers by enhancing the transport of unspliced viral RNA out of the nucleus (2). Finally, pLVX-shRNA1 also contains a central polypurine tract/central termination sequence element (cPPT/CTS). During target cell infection, thiselement creates a central DNA flap that increases nuclear import of the viral genome, resulting in improved vector integration and more efficient transduction (3). In addition to lentiviral elements, pLVX-shRNA1 contains a puromycin resistance gene (Puror)under the control of the murine phosphoglycerate kinase promoter (PPGK) for the selection of stable transductants. The vector also contains a pUC origin of replication and an E. coli ampicillin resistance gene (Ampr) for propagation and selection in bacteria.

质粒图谱



复溶方法

1. Centrifuge at 5,000×g for 5 min.
2. Carefully open the tube and add sterile water to dissolve the DNA: add 20 µl for 5 µg plasmid, and 100 µl for 100 µg plasmid.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than 5000×g.
5. Store the plasmid at -20 ℃.