
pLVX-IRES-tdTomato

载体编号 Vector ID	V012721
抗性 Antibiotic Resistance	Ampicillin
长度 Length	8892 bp
载体类型 Vector Type	Viral Expression & Packaging Vectors

产品信息

复制子 Replication Origin	ori
来源 Source/Author	Clontech
筛选标记 Selection Marker	tdTomato
拷贝数 Copy Number	High copy number
启动子 Promoter	CMV
克隆方法 Cloning Method	Enzyme Cut
表达方式 Expression Method	Constiutive, Stable
备注 Comment	pLVX-IRES-tdTomato is an HIV-1-based, lentiviral expression vector that allows the simultaneous expression of your protein of interest and tdTomato in virtually any mammalian cell type, including primary cells. tdTomato is a member of the family of fruit fluorescent proteins derived from the Discosoma sp. red fluorescent protein, DsRed (1). The vector expresses the two proteins from a bicistronic mRNA transcript, allowing tdTomato to be used as an indicator of transduction efficiency and a marker for selection by flow cytometry. Expression of the bicistronic transcript is driven by the constitutively active human cytomegalovirus immediate early promoter (PCMV IE) located just upstream of the MCS. An encephalomyocarditis virus (EMCV) internal ribosome entry site (IRES), positioned between the MCS and tdTomato, facilitates cap-independent translation of tdTomato from an internal start site at the IRES/tdTomato junction (1). pLVX-IRES-tdTomato contains all of the viral processing elements necessary for the production of replication-incompetent lentivirus, as well as elements to improve viral titer, transgene expression, and overall vector function. The woodchuck hepatitis virus posttranscriptional regulatory element (WPRE) promotes RNA processing events and enhances nuclear export of viral RNA (2), leading to increased viral titers from packaging cells. In addition, the vector includes a Rev-response element (RRE), which further increases viral titers by enhancing the transport of unspliced viral RNA out of the nucleus (3). Finally, pLVX-IRES-tdTomato also contains a central polypurine tract/central termination sequence element (cPPT/CTS). During target cell infection, this element creates a central DNA flap that increases nuclear import of the viral genome, resulting in improved vector integration and more efficient transduction (4). The vector also contains a pUC origin of replication and an E. coli ampicillin resistance gene (Ampr) for propagation and selection in bacteria.

质粒图谱



复溶方法

1. Centrifuge at 5,000×g for 5 min.
2. Carefully open the tube and add sterile water to dissolve the DNA: add 20 µl for 5 µg plasmid, and 100 µl for 100 µg plasmid.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than 5000×g.
5. Store the plasmid at -20 ℃.