Tcf/Lef:H2B-GFP

载体编号 Vector ID	V017328
抗性 Antibiotic Resistance	Kanamycin
长度 Length	5677 bp
载体类型 Vector Type	Protein expression
宿主 Host	Mammalian cells

产品信息

复制子 Replication Origin	ori
筛选标记 Selection Marker	Neo/G418
拷贝数 Copy Number	High copy number
启动子 Promoter	Hsp68
克隆方法 Cloning Method	Restriction Enzyme
5' 引物 5' Primer	5′ sequencing primerGF PGenotFW: A CAACAAGCGCTCGACCATCAC;
3' 引物 3' Primer	3′ sequencing primerGFPGenotRW: AGTCGATGCCCTTCAGCTCGAT
培养菌株 Growth Strain(s)	DH10B
培养温度 Growth Temperature	37℃
备注 Comment	Tcf/Lef:H2B-GFP places six TCF/Lef response elements plus an hsp68 minimal promoter upstream of nuclear H2B-GFP, reporting canonical Wnt/β-catenin activity at single-cell resolution in live mammalian cells and transgenic mice.

质粒图谱



复溶方法

1. Centrifuge at 5,000×g for 5 min.
2. Carefully open the tube and add sterile water to dissolve the DNA: add 20 µl for 5 µg plasmid, and 100 µl for 100 µg plasmid.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin to concentrate the liquid at the bottom. Speed is less than 5000×g.
5. Store the plasmid at -20 ℃.